The strong total rainbow connection of graphs*

Lin Chen, Xueliang Li, Jinfeng Liu

Center for Combinatorics and LPMC
Nankai University, Tianjin 300071, P.R. China

E-mail: chenlin1120120012@126.com, lxl@nankai.edu.cn, ljinfeng709@163.com

arXiv:1601.01063v2 [math.CO] 17 Mar 2016

Abstract

A graph is called *total colored* if both all its edges and all its vertices are colored. A path in a total colored graph is called *total rainbow* if all the edges and the internal vertices on the path have distinct colors. A graph is called *total rainbow connected* if any two vertices of the graph are connected by a total rainbow path. In this paper we introduce the concept of strong total rainbow connection of graphs. A graph is called *strongly total rainbow connected* if any two vertices of the graph are connected by a total rainbow geodesic, i.e., a path of length equal to the distance between the two vertices. For a connected graph G, the *strong total rainbow connection number*, denoted by $strc(G)$, is the minimum number of colors that are needed to make G strongly total rainbow connected. Among our results we state some simple observations about $strc(G)$ for a connected graph G. We also investigate the strong total rainbow connection numbers of some special graphs. Finally, for any pair of positive integers a and b with $b \geq a$, except for four cases, we can construct a connected graph G such that $trc(G) = a$ meanwhile $strc(G) = b$.

AMS Subject Classification 2010: 05C15, 05C40, 05C12.

1 Introduction

In this paper, all graphs under our consideration are finite, undirected and simple. For more notation and terminology that will be used in the sequel, we refer to [1], unless otherwise stated.

In 2008, Chartrand et al. [4] introduced the concept of rainbow connection. Let G be a nontrivial connected graph on which an edge-coloring $c : E(G) \to \{1, 2, \ldots, k\}(k \in \mathbb{N})$ is defined, where adjacent edges may be colored with the same color. For any two vertices u and v of G, a path in G connecting u and v is abbreviated as a uv-path. A uv-path P is a *rainbow uv-path* if no two edges of P are colored with the same color. The graph G is *rainbow connected* (with respect to c) if G contains a rainbow uv-path

*Supported by NSFC No.11371205 and 11531011.

for every two vertices u and v in it, and the coloring c is called a *rainbow coloring* of G. If k colors are used, then c is a *rainbow k-coloring*. For a connected graph G, the minimum k for which there exists a rainbow k-coloring of the edges of G is the *rainbow connection number* of G, denoted by $rc(G)$. The topic of rainbow connection is fairly interesting and numerous relevant papers have been published. In addition, the concept of strong rainbow connection was introduced by the same authors. For two vertices u and v of G, a *rainbow uv-geodesic* in G is a rainbow uv-path of length $d(u, v)$, where $d(u, v)$ is the distance between u and v (the length of a shortest uv-path in G). The graph G is *strongly rainbow connected* if G contains a rainbow uv-geodesic for every two vertices u and v of G. In this case, the coloring c is called a *strong rainbow coloring* of G. For a connected graph G, the minimum k for which there exists a coloring $c : E(G) \to \{1, 2, \ldots, k\}$ of the edges of G such that G is strongly rainbow connected is the *strong rainbow connection number* of G, denoted by $src(G)$. The investigation of $src(G)$ is more challenging than that of $rc(G)$, and very few papers have been obtained on it, for details see [4, 7, 11, 17].

As a natural counterpart of the rainbow connection, Krivelevich and Yuster proposed the concept of rainbow vertex-connection in [10]. A vertex colored graph G is *rainbow vertex-connected* if any two vertices of G are connected by a path whose internal vertices have different colors, and such a path is called a *vertex-rainbow path*. For a connected graph G, the *rainbow vertex-connection number* of G, denoted by $rvc(G)$, is the smallest number of colors needed to make G rainbow vertex-connected. Corresponding to the strong rainbow connection, Li et al. in [13] introduced the notion of strong rainbow vertex-connection. A vertex colored graph G is *strongly rainbow vertex-connected*, if for every pair of distinct vertices u and v, there exists a vertex-rainbow uv-geodesic. For a connected graph G, the minimum number k for which there exists a k-coloring of the vertices of G that results in a strongly rainbow vertex-connected graph is called the *strong rainbow vertex-connection number* of G, denoted by $srvc(G)$. For more results on rainbow vertex-connection, we refer to [14].

It was also shown that computing the rainbow connection number and rainbow vertex-connection number of an arbitrary graph is NP-hard [2, 3, 5, 6, 8, 12]. For more results on rainbow connections, we refer to the survey [15] and the book [16].

Subsequently, Liu et al. [18] proposed the concept of total rainbow connection. A *total colored graph* is a graph G such that both all edges and all vertices of G are colored. A total colored path is *total rainbow* if its edges and internal vertices have distinct colors. A total colored graph G is *total rainbow connected* if any two vertices of G are connected by a total rainbow path. For a connected graph G, the *total rainbow connection number* of G, denoted by $trc(G)$, is defined as the minimum number of colors required to make G total rainbow connected. Note the trivial fact that $trc(G) = 1$ if and only if G is a complete graph. Moreover, $2diam(G) - 1 \leq trc(G) \leq 2n - 3$ for any nontrivial connected graph G, where $diam(G)$ denotes the diameter of G. For more results on the total rainbow connection number, see [9, 18, 19].

Inspired by the concepts of total rainbow connection number $trc(G)$, strong rainbow connection number $src(G)$ and strong rainbow vertex-connection number $srvc(G)$, a natural idea is to introduce the concept of strong total rainbow connection number. Let G be a total colored graph. For any two vertices u and v of G, a *total rainbow uv-geodesic* in G is a total rainbow uv-path of length $d(u, v)$, where $d(u, v)$ is the distance between u and v. The graph G is *strongly total rainbow connected* if any two vertices of G are connected by a total rainbow geodesic. Such a coloring is called a strong total rainbow coloring of G. The *strong total rainbow connection number* of G, denoted by $strc(G)$, is defined to be the smallest number of colors needed to make G strongly total rainbow connected. Clearly, the trivial lower bound $strc(G) \geq 2diam(G) - 1$ is immediate from the definition, where G is any nontrivial connected graph and $diam(G)$ is the diameter of G. Further, this bound is tight as we will show that $strc(P_n) = 2n-3 = 2diam(P_n)-1$ for an n-vertex path P_n. Besides, by definition, $trc(G) \leq strc(G)$ holds for every connected graph G.

In this paper, we state some simple observations about $strc(G)$ for a nontrivial connected graph G. Then, we investigate the values of the strong total rainbow connection number of some special graphs. Lastly, for any pair of positive integers a and b with $b \geq a$, except for four cases, we construct a connected graph G such that $trc(G) = a$ meanwhile $strc(G) = b$.

2 Preliminaries

We in this section state some observations about $strc(G)$ for a nontrivial connected graph G, certain necessary lemmas are also listed.

For a nontrivial connected graph G, since $diam(G) = 1$ if and only if G is a complete graph, then the following proposition is immediate.

Proposition 1. *Let G be a nontrivial connected graph. Then $strc(G) = 1$ if and only if G is a complete graph; and $strc(G) \geq 3$ otherwise.*

Let G be a connected graph. For any strong total rainbow coloring of G, the colors of bridges and non-pendant vertices incident with bridges must be pairwise distinct. Hence, we have the following result.

Proposition 2. *Let G be a connected graph of order $n \geq 3$. Suppose that B is the set of all bridges in G, and C is the set of all non-pendant vertices incident with bridges. Denote $b = |B|$ and $c = |C|$, respectively. Then $strc(G) \geq b + c$.*

Apparently, for a nontrivial connected graph G, according to the definition of $strc(G)$, $strc(G) \geq \max\{srvc(G), src(G)\}$ holds trivially. Additionally, the following associative upper bound is obvious and we omit the proof.

Proposition 3. *Let G be a connected graph with n vertices and p pendant vertices. Then $strc(G) \leq src(G) + n - p$.*

In [18], the authors derived an upper bound on the total rainbow connection number, which will be helpful in the next section.

Lemma 1 ([18])**.** *Let G be a connected graph on n vertices, with q vertices having degree at least 2. Then, $trc(G) \leq n - 1 + q$, with equality if and only if G is a tree.*

Let C_n denote the cycle of order n. The total rainbow connection number of C_n for $n \geq 3$ is established in [18].

Lemma 2 ([18])**.** *For $3 \leq n \leq 12$, the values of $trc(C_n)$ are given in the following table.*

n	3	4	5	6	7	8	9	10	11	12
$trc(C_n)$	1	3	3	5	6	7	8	9	11	11

For $n \geq 13$, we have $trc(C_n) = n$.

The *wheel* W_n is the graph obtained from the cycle C_n by joining a new vertex v to every vertex of C_n. The vertex v is the *centre* of W_n. $K_{m,n}$ denotes the complete bipartite graph with bipartites of sizes m and n. In [18], the authors determined $trc(W_n)$ for $n \geq 3$ and $trc(K_{m,n})$ for $2 \leq m \leq n$.

Lemma 3 ([18])**.** *$trc(W_3) = 1$, $trc(W_n) = 3$ for $4 \leq n \leq 6$, $trc(W_n) = 4$ for $7 \leq n \leq 9$, and $trc(W_n) = 5$ for $n \geq 10$.*

Lemma 4 ([18])**.** *For $2 \leq m \leq n$, we have $trc(K_{m,n}) = \min(\lceil \sqrt[m]{n} \rceil + 1, 7)$.*

3 Strong total rainbow connection numbers of some graphs

In this section, we study the strong total rainbow connection numbers for some specific graphs. We begin with trees of order n.

Theorem 1. *Let T be a nontrivial tree of order n with p pendant vertices. Then $strc(T) = 2n - p - 1$.*

Proof. From Lemma 1 we know that $trc(T) = 2n - p - 1$. Thus $strc(T) \geq trc(T) = 2n - p - 1$.

On the other hand, for any total rainbow coloring c of T with $trc(T) = 2n - p - 1$ colors and arbitrary two vertices u and v of T, the total rainbow path between u and

v is also the total rainbow geodesic between u and v since u and v are connected by exactly one path. Therefore, c is also a strong total rainbow coloring of T. That is, $strc(T) \leq 2n - p - 1$. Hence $strc(T) = 2n - p - 1$. ∎

Remark: By Theorem 1, we have $trc(T) = strc(T)$ for any tree T. Hence, for any integer $a \geq 5$, we can construct a tree T of order n ($\frac{a+3}{2} \leq n \leq a$) with $2n - a - 1$ pendant vertices satisfying $trc(T) = strc(T) = a$.

It is well known that every nontrivial tree has at least two pendant vertices and every nontrivial tree with $n \geq 3$ has at most $n - 1$ pendant vertices. Thus for a nontrivial tree T of order $n \geq 3$, we have $n \leq strc(T) \leq 2n - 3$. In addition, both the lower bound and the upper bound can be reached by the n-vertex star S_n and the n-vertex path P_n. Particularly, $strc(P_n) = 2n - 3 = 2diam(P_n) - 1$, as mentioned in our introduction.

By taking full advantage of the fact that $strc(G) \geq trc(G)$ and the proof of Lemma 2 (see the proof of Theorem 2 in [18]), one can easily check that $strc(C_3) = 1$, $strc(C_4) = 3$, $strc(C_5) = 3$, $strc(C_6) = 5$, $strc(C_7) = 6$, $strc(C_8) = 7$, $strc(C_9) = 8$, $strc(C_{10}) = 9$, $strc(C_{11}) = 11$, $strc(C_{12}) = 11$. We aim at determining $strc(C_n)$ for $n \geq 13$ in the following theorem.

Theorem 2. *For $n \geq 13$, we have $strc(C_n) = n$.*

Proof. First, since $strc(G) \geq trc(G)$ for any connected graph G. The lower bound $strc(C_n) \geq n$ is easy to get by Lemma 2.

Then, we prove the upper bound. By the definition of strong total rainbow connection number, it suffices to find a strong total rainbow coloring using n colors for C_n. Let $C_n = v_0v_1 \cdots v_{n-1}v_0$. We discuss the following two cases.

Case 1 If n is even.

Then, color the edges v_0v_1, v_1v_2, ..., $v_{\frac{n}{2}-1}v_{\frac{n}{2}}$, $v_{\frac{n}{2}}v_{\frac{n}{2}+1}$, $v_{\frac{n}{2}+1}v_{\frac{n}{2}+2}$, ..., $v_{n-1}v_0$ with colors 0, 1, ..., $\frac{n}{2} - 1$, 0, 1, ..., $\frac{n}{2} - 1$, respectively. Meanwhile, gradually color the vertices v_0, v_1, ..., $v_{\frac{n}{2}-1}$, $v_{\frac{n}{2}}$, ..., v_{n-1} with colors $\frac{n}{2}$, $\frac{n}{2} + 1$, ..., $n - 1$, $\frac{n}{2}$, ..., $n - 1$. Whereupon, a total coloring using n colors is assigned to C_n. Under this coloring, any path with length no more than $\frac{n}{2}$ is a total rainbow path. Since $diam(C_n) = \frac{n}{2}$, then for any two vertices v_i, v_j of C_n, there is a total rainbow v_iv_j-geodesic. Hence, C_n is strongly total rainbow connected. That is, $strc(C_n) \leq n$.

Case 2 If n is odd.

Then, color the edges v_0v_1, v_1v_2, ..., $v_{\frac{n-3}{2}}v_{\frac{n-1}{2}}$, $v_{\frac{n+1}{2}}v_{\frac{n+3}{2}}$, $v_{\frac{n+3}{2}}v_{\frac{n+5}{2}}$, ..., $v_{n-1}v_0$ with colors 0, 1, ..., $\frac{n-3}{2}$, 0, 1, ..., $\frac{n-3}{2}$ stepwise. The remaining edge $v_{\frac{n-1}{2}}v_{\frac{n+1}{2}}$ is assigned the color $\frac{n-1}{2}$. Next, color the vertices v_1, v_2, ..., $v_{\frac{n-1}{2}}$, $v_{\frac{n+1}{2}}$, ..., v_{n-1} with colors $n-1$, $n-2$, ..., $\frac{n+1}{2}$, $n-1$, ..., $\frac{n+1}{2}$. Further, assign the color $\frac{n-1}{2}$ to the vertex v_0. Now we have a total coloring of C_n with n colors. Similarly, under this coloring, one can check

that for any pair of vertices v_i and v_j $(i \neq j)$, there exists a total rainbow v_iv_j-geodesic in C_n. Accordingly, $strc(C_n) \leq n$.

In summary, we finally arrive at $strc(C_n) = n$ for $n \geq 13$. ∎

We give the following theorem to study the strong total rainbow connection number of W_n, whose proof is partially based on the fact that $strc(W_n) \geq trc(W_n)$.

Theorem 3. *Let W_n be a wheel with $n \geq 3$. Then*

$$\begin{cases} strc(W_3) = 1, \\ strc(W_n) = \lceil \frac{n}{3} \rceil + 1 \quad \textit{for } n \geq 4. \end{cases}$$

Proof. Let v be the centre of W_n, and v_1, v_2, ..., v_n be the vertices of W_n in the cycle C_n.

(i) Since W_3 is precisely the complete graph K_4, then obviously $strc(W_3) = 1$.

(ii) It follows from Lemma 3 that $strc(W_n) \geq trc(W_n) = 3$ when $4 \leq n \leq 6$. On the other hand, when $4 \leq n \leq 6$, we define a total coloring f_n with 3 colours for W_n as the same as Liu et al. [18]. Namely, let $f_n(vv_i) = \lceil \frac{i}{3} \rceil$, $f_n(v_iv_{i+1}) = 1$ if $i \equiv 1 \pmod 3$, $f_n(v_iv_{i+1}) = 2$ if $i \equiv 2 \pmod 3$, $f_n(v_iv_{i+1}) = 3$ if $i \equiv 0 \pmod 3$ and $f_n(v) = f_n(v_i) = 3$ for $i = 1, 2, \ldots, n$. Note that $v_{n+1} = v_1$. Then, under this coloring, W_n is strongly total rainbow connected. Hence $strc(W_n) \leq 3$. Therefore, we get that $strc(W_n) = 3 = \lceil \frac{n}{3} \rceil + 1$ for $4 \leq n \leq 6$.

(iii) What remains to be shown is that $strc(W_n) = \lceil \frac{n}{3} \rceil + 1$ for $n \geq 7$. Firstly, we prove the lower bound, that is, $strc(W_n) \geq \lceil \frac{n}{3} \rceil + 1$. Assume to the contrary, that $strc(W_n) \leq \lceil \frac{n}{3} \rceil$. Let f'_n be a strong total rainbow coloring of W_n with $strc(W_n)$ colors. Since $n \geq 7$, then for each vertex $v_i (1 \leq i \leq n)$, there exists at least one vertex v_j with $j \in \{1, 2, \ldots, n\}, j \neq i$ such that the unique v_iv_j-geodesic of length 2 passes the centre v. Thus, $f'_n(v) \neq f'_n(vv_i)$ for $i = 1, 2, \ldots, n$. Therefore, the n edges $vv_i (i = 1, 2, \ldots, n)$ use at most $strc(W_n) - 1 \leq \lceil \frac{n}{3} \rceil - 1 < \frac{n}{3}$ different colors. Which deduces that there exist at least four different edges, say vv_i, vv_j, vv_k, vv_l, such that $f'_n(vv_i) = f'_n(vv_j) = f'_n(vv_k) = f'_n(vv_l)$. Again, since $n \geq 7$, there exist at least two vertices, without loss of generality, say v_i and v_j, such that the unique v_iv_j-geodesic is precisely the path v_ivv_j. So, there are no total rainbow v_iv_j-geodesics as $f'_n(vv_i) = f'_n(vv_j)$, a contradiction. Consequently, $strc(W_n) \geq \lceil \frac{n}{3} \rceil + 1$.

Now we prove the upper bound by giving W_n a strong total rainbow coloring g_n with colors from $\{1, 2, \ldots, \lceil \frac{n}{3} \rceil + 1\}$, as follows. Let $g_n(vv_i) = \lceil \frac{i}{3} \rceil$ for $i = 1, 2, \ldots, n$; $g_n(v) = \lceil \frac{n}{3} \rceil + 1$; $g_n(v_iv_{i+1}) = 1$ if $i \equiv 1 \pmod 3$; $g_n(v_iv_{i+1}) = 2$ if $i \equiv 2 \pmod 3$; $g_n(v_iv_{i+1}) = 3$ if $i \equiv 0 \pmod 3$; $g_n(v_{3j+2}) = 4$ for $j = 0, 1, \ldots, \lfloor \frac{n-2}{3} \rfloor$. At last, color the remaining vertices arbitrarily with colors in $\{1, 2, \ldots, \lceil \frac{n}{3} \rceil + 1\}$. Notice that here we stipulate $v_{n+1} = v_1$ and since $n \geq 7$, we have at least $\lceil \frac{7}{3} \rceil + 1 = 4$ different colors available. One can check that this total coloring g_n is indeed a strong total rainbow coloring of W_n. Accordingly, by definition, $strc(W_n) \leq \lceil \frac{n}{3} \rceil + 1$.

In conclusion, we have established that $strc(W_n) = \lceil \frac{n}{3} \rceil + 1$ for $n \geq 7$, and then the theorem is proved. ∎

By Proposition 1 and Theorem 1, we know that $strc(K_{1,1}) = 1$ and $strc(K_{1,n}) = n+1$ for $n \geq 2$. In the sequel, we will determine $strc(K_{m,n})$ for $2 \leq m \leq n$.

Theorem 4. *For $2 \leq m \leq n$, we have $\lceil \sqrt[m]{n} \rceil \leq strc(K_{m,n}) \leq \lceil \sqrt[m]{n} \rceil + 1$.*

Proof. Let the bipartition of $K_{m,n}$ be U and V, where $U = \{u_1, \ldots, u_m\}$ and $V = \{v_1, \ldots, v_n\}$. We first show the lower bound by contradiction. Assume that $strc(K_{m,n}) < \lceil \sqrt[m]{n} \rceil$, that is, $strc(K_{m,n}) \leq \lceil \sqrt[m]{n} \rceil - 1$. Then, there exists a strong total rainbow coloring, say g, using colors from $\{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil - 1\}$ such that $K_{m,n}$ is strongly total rainbow connected. Assign a vector $z_i = (z_{i1}, z_{i2}, \ldots, z_{im})$ of length m to every vertex $v_i (i = 1, 2, \ldots, n)$ satisfying $z_{ij} = g(v_i u_j) (j = 1, 2, \ldots, m)$. Since $g(v_i u_j) \in \{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil - 1\}$, there are at most $(\lceil \sqrt[m]{n} \rceil - 1)^m$ distinct vectors. Clearly, $(\lceil \sqrt[m]{n} \rceil - 1)^m < n$. Therefore, there exist at least two vertices v_{k_1} and v_{k_2} such that $z_{k_1} = z_{k_2}$. Namely, $g(v_{k_1} u_j) = g(v_{k_2} u_j)$ for each $j \in \{1, 2, \ldots, m\}$. Since any $v_{k_1} v_{k_2}$-geodesic in $K_{m,n}$ must contain some vertex of U, then there exist no total rainbow geodesics between v_{k_1} and v_{k_2}, which contradicts that g is a strong total rainbow coloring. Hence, $strc(K_{m,n}) \geq \lceil \sqrt[m]{n} \rceil$.

Now we begin to show the upper bound. We construct a strong total rainbow coloring f for $K_{m,n}$ by using colors from $\{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil + 1\}$ as follows. Assign to the vertices of V distinct vectors of length m with entries from $\{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil\}$. Since $(\lceil \sqrt[m]{n} \rceil)^m \geq n = |V|$, we can make sure that any pair of vertices of V are assigned with different vectors. Simultaneously, since $m \leq n$, we can stipulate that the m vectors $(2, 1, 1, \ldots, 1)$, $(1, 2, 1, \ldots, 1)$, $\ldots$, $(1, 1, \ldots, 2, 1)$, $(1, 1, \ldots, 1, 2)$ are all present. For $v_i \in V$, let $z'_i = (z'_{i1}, z'_{i2}, \ldots, z'_{im})$ denote the vector assigned to v_i. Then, for $u_j \in U$ and $v_i \in V$, let $f(u_j v_i) = z'_{ij}$ $(j = 1, 2, \ldots, m; i = 1, 2, \ldots, n)$. Further, let $f(w) = \lceil \sqrt[m]{n} \rceil + 1$ for all $w \in V(K_{m,n})$. In such a way, we define a total coloring for $K_{m,n}$. Under this coloring, for u_{j_1}, $u_{j_2} \in U$, the path $u_{j_1} z u_{j_2}$ is a total rainbow geodesic between u_{j_1} and u_{j_2}, where $z \in V$ is the vertex assigned the vector $z' = (1, \ldots, 1, 2, 1, \ldots, 1)$, with 2 in the j_1-th position. For v_{k_1}, $v_{k_2} \in V$, since $z'_{k_1} \neq z'_{k_2}$, there exists at least one index $j \in \{1, 2, \ldots, m\}$ such that $z'_{k_1 j} \neq z'_{k_2 j}$. It follows that there exists at least one vertex $u_j \in U$ such that $f(u_j v_{k_1}) \neq f(u_j v_{k_2})$. Hence, the path $v_{k_1} u_j v_{k_2}$ is a total rainbow geodesic between v_{k_1} and v_{k_2}. While for any $u \in U$ and any $v \in V$, there obviously exists a total rainbow uv-geodesic. Therefore, for any two vertices of $K_{m,n}$, there exists at least one total rainbow geodesic connecting them, namely, $K_{m,n}$ is strongly total rainbow connected under the coloring f. Hence, $strc(K_{m,n}) \leq \lceil \sqrt[m]{n} \rceil + 1$.

To sum up, we arrive at $\lceil \sqrt[m]{n} \rceil \leq strc(K_{m,n}) \leq \lceil \sqrt[m]{n} \rceil + 1$. ∎

Actually, in Theorem 4, if $strc(K_{m,n}) = \lceil \sqrt[m]{n} \rceil$, then for any strong total rainbow coloring g of $K_{m,n}$ with colors from $\{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil\}$, we conjecture that there exists at least one vertex $w \in V(K_{m,n})$ such that $g(w)$ is different from all the colors of the

edges in $K_{m,n}$, that is, the edges of $K_{m,n}$ use at most $\lceil \sqrt[m]{n} \rceil - 1$ colors, which will deduce a contradiction. Hence, we pose the following conjecture.

Conjecture 1. *If $2 \leq m \leq n$, then $strc(K_{m,n}) = \lceil \sqrt[m]{n} \rceil + 1$.*

It's worth mentioning that according to the proof of Theorem 4 in [18], we know that $trc(K_{m,n}) = \lceil \sqrt[m]{n} \rceil + 1$ for $m \leq n \leq 6^m$. Hence, $strc(K_{m,n}) \geq trc(K_{m,n}) = \lceil \sqrt[m]{n} \rceil + 1$ if $m \leq n \leq 6^m$, which means that the conjecture above holds for $2 \leq m \leq n \leq 6^m$.

At last, we discuss the case when G is a complete multipartite graph $K_{n_1,\ldots,n_t}$. Let M be a matching. The two ends of each edge in M are said to be matched under M, and each vertex incident with an edge in M is said to be covered by M. Moreover, a matching M is called a k-matching if the size of M is k.

Theorem 5. *Let $t \geq 3$, $1 \leq n_1 \leq \cdots \leq n_t$, $m = \sum_{i=1}^{t-1} n_i$ and $n = n_t$. Then,*

$$\begin{cases} strc(K_{n_1,\ldots,n_t}) = 1 & \text{if } n = 1; \\ strc(K_{n_1,\ldots,n_t}) = 3 & \text{if } n \geq 2 \text{ and } m > n; \\ \lceil \sqrt[m]{n} \rceil \leq strc(K_{n_1,\ldots,n_t}) \leq \lceil \sqrt[m]{n} \rceil + 1 & \text{if } n \geq 2 \text{ and } m \leq n. \end{cases}$$

Proof. For convenience, we write G instead of $K_{n_1,\ldots,n_t}$. Let V_i be the i-th partite with n_i vertices for $1 \leq i \leq t$. Set $U = \bigcup_{i=1}^{t-1} V_i$.

(i) If $n = 1$, then $G = K_{n_1,\ldots,n_t} = K_{1,\ldots,1} = K_t$ is a complete graph. Hence, clearly $strc(G) = strc(K_t) = 1$.

(ii) If $n \geq 2$ and $m > n$, then G is not complete, which implies that $strc(G) \geq 3$. Since $|U| = \sum_{i=1}^{t-1} |V_i| = \sum_{i=1}^{t-1} n_i = m > n$, there exists an n-matching E_0 between U and V_t. Let $U_0 = \{u \in U : u \text{ is covered by } E_0\}$, $U_i = V_i \cap U_0$ and $E_0|_{U_i} = \{e = uv \in E_0 : u \in U_i \text{ and } v \in V_t\}$. For each i with $1 \leq i \leq t-1$, since $n_i \leq n_t$, there is a $|V_i \setminus U_i|$-matching E_i between $V_i \setminus U_i$ and V_t such that E_i and $E_0|_{U_i}$ are vertex-disjoint. In order to prove $strc(G) \leq 3$ in this case, we give G such a total coloring c using 3 colors, as defined as follows. Let $c(e) = 1$ for $e \in E_0 \cup E_1 \cdots \cup E_{t-1}$; $c(v) = 2$ for all $v \in G$; color the remaining edges with color 3. We will show that c is a strong total rainbow coloring of G. Let u and v be any two different vertices of G. If u and v are in different partites, then there exists a total rainbow geodesic connecting u and v, obviously. If u and v are in the same partite, then according to the positions of u and v, the discussion is divided into four aspects regardless of the symmetric cases.

Case 1 $u, v \in V_t$.

Then, there exists some vertex $w \in U$ such that $wv \in E_0$ while $wu \notin E_0 \cup E_1 \cdots \cup E_{t-1}$. By the definition of the coloring c, we know that $c(wv) = 1$, $c(w) = 2$, $c(wu) = 3$. Hence, uwv is a total rainbow geodesic between u and v.

Case 2 $u, v \in U_i$ for some $i \in \{1, 2, \ldots, t-1\}$.

Then, there exists some vertex $x \in V_t$ such that $vx \in E_0$ while $ux \notin E_0 \cup E_1 \cdots \cup E_{t-1}$. Hence, $c(vx) = 1$, $c(x) = 2$, $c(ux) = 3$. Accordingly, uxv is a total rainbow geodesic connecting u and v.
Case 3 $u \in U_i$, $v \in V_i \setminus U_i$ for some $i \in \{1, 2, \ldots, t-1\}$.
Case 4 $u, v \in V_i \setminus U_i$ for some $i \in \{1, 2, \ldots, t-1\}$.
The arguments of Cases 3 and 4 are similar to that of Case 2, we will not elaborate.

Based on the above analysis, we have illustrated the existence of a total rainbow uv-geodesic for any two vertices u and v in G. Thus, c is a strong total rainbow coloring of G with 3 colors. It follows that $strc(G) \leq 3$. Therefore, $strc(G) = 3$ if $n \geq 2$ and $m > n$.

(iii) If $n \geq 2$ and $m \leq n$, we first prove the lower bound. Consider the edges between U and V_t. For any two vertices u, $v \in V_t$, any uv-geodesic must contain a vertex of U. Similar to the argument for the lower bound of strong total rainbow connection number for complete bipartite graphs, we can get $strc(G) \geq \lceil \sqrt[m]{n} \rceil$.
Then, we prove the upper bound by defining a strong total rainbow coloring c for G with $\lceil \sqrt[m]{n} \rceil + 1$ colors. Let $U = \{v_1, v_2, \ldots, v_m\}$. Assign to the vertices of V_t distinct vectors of length m with entries from $\{1, 2, \ldots, \lceil \sqrt[m]{n} \rceil\}$. This is possible since $(\lceil \sqrt[m]{n} \rceil)^m \geq n = |V_t|$. In addition, since $m \leq n$, the vectors $(2, 1, 1, \ldots, 1)$, $(1, 2, 1, \ldots, 1)$, $\ldots$, $(1, 1, \ldots, 2, 1)$, $(1, 1, \ldots, 1, 2)$ can all be present. For $v \in V_t$, let $v' = (v'_1, v'_2, \ldots, v'_m)$ denote the vector assigned to v. Then, for $v_i \in U$ and $v \in V_t$, let $c(vv_i) = v'_i$. For the remaining edges, we color them with color $\lceil \sqrt[m]{n} \rceil + 1$. Finally, let $c(v) = \lceil \sqrt[m]{n} \rceil + 1$ for all $v \in G$. One can check that such a coloring c is actually a strong total rainbow coloring of G with $\lceil \sqrt[m]{n} \rceil + 1$ colors (the proof is the same as that for Theorem 4). Hence, $strc(G) \leq \lceil \sqrt[m]{n} \rceil + 1$. Consequently, $\lceil \sqrt[m]{n} \rceil \leq strc(G) \leq \lceil \sqrt[m]{n} \rceil + 1$ if $n \geq 2$ and $m \leq n$.

The proof is thus complete. ∎

4 The (strong) total rainbow connection numbers with prescribed values

We have seen that $strc(G) \geq trc(G)$ holds for every connected graph G. In addition, we also find graphs for which the equality holds. For example, $strc(C_n) = trc(C_n) = n$ if $n \geq 13$. However, we want to know what is the difference between these two parameters. In this section, we aim to find a connected graph G for every pair of positive integers a, b with $b \geq a$, except for four cases, such that $trc(G) = a$ while $strc(G) = b$.

In order to prove the main result, we need some assistant lemmas.

Lemma 5. *Let G be a connected graph and $e = uv$ be a pendant edge of G with pendant vertex u. Then, $trc(G - u) \leq trc(G)$.*

Proof. By definition, for a given total rainbow coloring of G with $trc(G)$ colors, there is a total rainbow path between each pair of vertices of G. Since e is a pendant edge of G, every total rainbow path connecting two vertices of $G-u$ misses e. Therefore, there is still a total rainbow path between any two vertices of $G-u$, which implies that $trc(G-u) \leq trc(G)$. ∎

Remark 1: Similarly, we can show that the result of Lemma 5 is also true for the version of strong total rainbow connection number, that is, we also have $strc(G-u) \leq strc(G)$.

Before we give the following lemmas, a new graph is constructed. Denote by W_n^r ($r \geq 1$) the graph obtained from a wheel W_n by attaching r pendant edges at the center v of W_n. The r pendant vertices are denoted by $u_1, u_2, \ldots, u_r$ and the n vertices in the cycle C_n of W_n are denoted by $v_1, v_2, \ldots, v_n$. We will investigate $trc(W_n^r)$ and $strc(W_n^r)$, respectively.

Lemma 6. *For $n \geq 10$, we have that $trc(W_n^r) = 5$ for $1 \leq r \leq 4$, and $trc(W_n^r) = r+1$ for $r \geq 5$.*

Proof. According to Lemmas 3 and 5, we know that $trc(W_n^r) \geq trc(W_n) = 5$ when $n \geq 10$. For W_n^4, we give it a total coloring g as follows. Let $g(vv_i) = 1$ if $i \equiv 1 \pmod 2$, $g(vv_i) = 2$ if $i \equiv 0 \pmod 2$, $g(v_i) = 3$ for $i = 1, 2, \ldots, n$, $g(v_iv_{i+1}) = 4$ for $i = 1, 2, \ldots, n$, where $v_{n+1} = v_1$, $g(v) = g(u_i) = 5$ and $g(vu_i) = i$ for $i = 1, 2, 3, 4$. Then, one can check that g is indeed a total rainbow coloring of W_n^4 using 5 colors. Hence, $trc(W_n^4) \leq 5$. It follows that $trc(W_n^4) = 5$. In addition, for $1 \leq r \leq 3$, we have that $trc(W_n^r) \leq trc(W_n^4) = 5$ by Lemma 5. Thus, $trc(W_n^r) = 5$ for $r = 1, 2, 3$.

Now we discuss the case $r \geq 5$. Let u_i and u_j ($i \neq j$, $i, j = 1, 2, \ldots, r$) be any two pendant vertices in W_n^r. In order to get a total rainbow path connecting u_i and u_j, vu_i and vu_j must have different colors, which means that the r pendant edges must be assigned r different colors. Furthermore, any total rainbow path connecting two pendant vertices passes the vertex v. Hence, the color of v is different from the color of vu_i for each $i \in \{1, 2, \ldots, r\}$. In summary, to get a total rainbow coloring of W_n^r, at least $r+1$ colors are required, namely, $trc(W_n^r) \geq r+1$. We in the following construct a total coloring f for W_n^r by using colors from $\{1, 2, \ldots, r+1\}$.

Let $f(vv_i) = 1$ if $i \equiv 1 \pmod 2$, $f(vv_i) = 2$ if $i \equiv 0 \pmod 2$, $f(v_i) = 3$ and $f(v_iv_{i+1}) = 4$ for $i = 1, 2, \ldots, n$, similarly, $v_{n+1} = v_1$, $f(v) = 5$, $f(u_i) = 5$ for $i = 1, 2, \ldots, r$, $f(vu_i) = i$ for $i = 1, 2, 3, 4$ and $f(vu_i) = i+1$ for $i = 5, 6, \ldots, r$. Under this coloring, we can find a total rainbow path for any two vertices of W_n^r. Hence, $trc(W_n^r) \leq r+1$. Accordingly, $trc(W_n^r) = r+1$ for $r \geq 5$. The lemma is thus proved. ∎

Lemma 7. *For $n \geq 10$, we have $strc(W_n^r) = \lceil \frac{n}{3} \rceil + r + 1$.*

Proof. At first, we prove the lower bound. For any two pendant vertices u_i and u_j in W_n^r, where $i \neq j$, $i, j = 1, 2, \ldots, r$, in order to obtain a total rainbow geodesic

connecting them, the colors of vu_i, v and vu_j are all distinct, which deduces that there should be $r+1$ distinct colors for $v, vu_1, vu_2, \ldots, vu_r$. Without loss of generality, let these $r+1$ colors be $1, 2, \ldots, r+1$, respectively. Moreover, let v_k $(k = 1, 2, \ldots, n)$ and u_i $(i = 1, 2, \ldots, r)$ be any two vertices in W_n^r. Since $d(v_k, u_i) = 2$, to make sure that there exists a total rainbow geodesic between v_k and u_i, the colors of vv_k, v and vu_i are all distinct. Hence, the color of vv_k is different from the colors in $\{1, 2, \ldots, r+1\}$, i.e., new colors should be used for the edges vv_k $(k = 1, 2, \ldots, n)$. We claim that among these edges, there are at most three edges assigned the same color.

Assume, to the contrary, that there exist at least four edges having the same color, say vv_{k_1}, vv_{k_2}, vv_{k_3}, vv_{k_4}. Since $n \geq 10$, then among the four vertices v_{k_1}, v_{k_2}, v_{k_3} and v_{k_4}, there are at least two of them having distance greater than 2 in C_n. Hence, there exists no total rainbow geodesic between such two vertices, a contradiction. Therefore, the n edges vv_k $(k = 1, 2, \ldots, n)$ use at least $\lceil \frac{n}{3} \rceil$ distinct colors. As a consequence, $strc(W_n^r) \geq \lceil \frac{n}{3} \rceil + r + 1$.

We now prove the upper bound by defining a strong total rainbow coloring c for W_n^r as follows: $c(vu_i) = i$ for $i = 1, 2, \ldots, r$; $c(v) = r+1$; $c(vv_k) = c(vv_{k+1}) = c(vv_{k+2}) = r + 1 + \lceil \frac{k}{3} \rceil$ for $k = 1, 4, 7, \ldots, 3(\lceil \frac{n}{3} \rceil - 1) + 1$; $c(v_k v_{k+1}) = r+1$ if $k \equiv 1 \pmod 3$, $c(v_k v_{k+1}) = r+2$ if $k \equiv 2 \pmod 3$, $c(v_k v_{k+1}) = r+3$ if $k \equiv 0 \pmod 3$, where $k = 1, 2, \ldots, n$ and $v_{n+1} = v_1$; $c(v_{3j+2}) = r+4$ for $j = 0, 1, \ldots, \lfloor \frac{n-2}{3} \rfloor$; color the remaining vertices with arbitrary colors from $\{1, 2, \ldots, \lceil \frac{n}{3} \rceil + r + 1\}$. Under this coloring, one can check that any two vertices of W_n^r have a total rainbow geodesic connecting them. Hence, this coloring is really a strong total rainbow coloring of W_n^r with $\lceil \frac{n}{3} \rceil + r + 1$ colors. Consequently, $strc(W_n^r) \leq \lceil \frac{n}{3} \rceil + r + 1$.

Combining with the lower and upper bounds, we finally arrive at $strc(W_n^r) = \lceil \frac{n}{3} \rceil + r + 1$. ∎

With the aid of Lemmas 6 and 7, we can arrive at the following result.

Theorem 6. *Let a and b be any two integers with $a = 5$ and $b \geq 6$ or $a \geq 6$ and $b \geq a+4$. Then, there exists a connected graph G such that $trc(G) = a$ and $strc(G) = b$.*

Proof. In the following, we shall apply Lemmas 6 and 7 repeatedly, for which we will no longer state.

If $a = 5$ and $6 \leq b \leq 9$, let $G = W_{12}^{b-5}$. Since $6 \leq b \leq 9$, then $1 \leq b - 5 \leq 4$. Hence, $trc(G) = trc(W_{12}^{b-5}) = 5 = a$, and $strc(G) = strc(W_{12}^{b-5}) = \lceil \frac{12}{3} \rceil + b - 5 + 1 = b$.

If $a = 5$ and $b \geq 10$, let $G = W_{3b-15}^4$, where $3b - 15 > 10$. Then, $trc(G) = trc(W_{3b-15}^4) = 5 = a$ and $strc(G) = strc(W_{3b-15}^4) = \lceil \frac{3b-15}{3} \rceil + 4 + 1 = b$.

If $a \geq 6$ and $b \geq a + 4$. Let $G = W_{3b-3a}^{a-1}$, where $a - 1 \geq 5$ and $3b - 3a > 10$. Then, $trc(G) = trc(W_{3b-3a}^{a-1}) = a - 1 + 1 = a$, meanwhile $strc(G) = strc(W_{3b-3a}^{a-1}) = \lceil \frac{3b-3a}{3} \rceil + a - 1 + 1 = b$.

The proof of this theorem is thus complete. ∎

As we all know, $strc(G) \geq trc(G)$ for any connected graph G. In the theorem above, we construct a connected graph G with $trc(G) = a$ and $strc(G) = b$, where $a \geq 6$, $b \geq a + 4$. Moreover, $strc(T) = trc(T)$ for any tree T. Then, whether exists a connected graph G such that $trc(G) < strc(G) \leq trc(G) + 3$? In order to solve this problem, we introduce a new kind of graphs.

Let W_t be a wheel, as defined before. Denote by $W_t^{r,s}$ the graph obtained from W_t by joining r and s pendant vertices to two adjacent vertices of C_t in W_t, respectively, where $r \geq 1$, $s \geq 1$. Let v be the centre of W_t and the vertices of the cycle C_t be $v_1, v_2, \ldots, v_t$. Without loss of generality, suppose that the r pendant vertices $u_1, u_2, \ldots, u_r$ and s pendant vertices $w_1, w_2, \ldots, w_s$ are joined to v_1 and v_2, respectively. According to the structure of $W_t^{r,s}$, we can arrive at the following result easily.

Claim 1. *Let r and s be two positive integers. Then, $trc(W_t^{r,s}) \geq r + s + 2$.*

Proof of Claim 1. Let c be any total rainbow coloring of $W_t^{r,s}$. For any two vertices u_i and u_j, $i \neq j$ and $i, j \in \{1, 2, \ldots, r\}$, the unique total rainbow u_iu_j-path is $u_iv_1u_j$, and thus, the colors $c(v_1)$, $c(v_1u_i)$ and $c(v_1u_j)$ are different. Similarly, we can obtain that the colors $c(v_2)$, $c(v_2w_k)$ and $c(v_2w_l)$ are different, where $k \neq l$ and $k, l \in \{1, 2, \ldots, s\}$. In addition, the total rainbow u_iw_k-path, $i \in \{1, 2, \ldots, r\}$, $k \in \{1, 2, \ldots, s\}$, must contain the edges v_1u_i, v_2w_k and v_1, v_2 as internal vertices, and hence, the colors $c(v_1)$, $c(v_2)$, $c(v_1u_i)$ and $c(v_2w_k)$ are distinct.

In summary, the colors $c(v_1)$, $c(v_2)$, $c(v_1u_i)(i = 1, 2, \ldots, r)$ and $c(v_2w_k)(k = 1, 2, \ldots, s)$ are pairwise different in any total rainbow coloring c, which deduces that $trc(W_t^{r,s}) \geq r + s + 2$.

Lemma 8. *Let r be an integer. Then, $trc(W_4^{r,1}) = r + 3$ for $r \geq 4$; $trc(W_7^{r,1}) = r + 3$ for $r \geq 5$; $trc(W_{10}^{r,1}) = r + 3$ for $r \geq 8$.*

Proof. By Claim 1, $trc(W_t^{r,1}) \geq r + 1 + 2 = r + 3$, which is a lower bound for $trc(W_t^{r,1})$. Then, it suffices to show $trc(W_t^{r,1}) \leq r + 3$ when $t = 4$, 7, 10, respectively, with given range of r.

If $t = 4$ and $r \geq 4$. Define a total coloring c_1 of $W_4^{r,1}$ using colors $1, 2, \ldots, r + 3$ as follows:
$c_1(v_1u_i) = i$ for $1 \leq i \leq r$; $c_1(v_1v_2) = c_1(v_3v_4) = c_1(vv_4) = 1$; $c_1(vv_1) = c_1(v_2v_3) = 2$; $c_1(v) = 3$; $c_1(vv_2) = 4$; $c_1(v_2w_1) = c_1(vv_3) = c_1(v_4) = r + 1$; $c_1(v_1) = r + 2$; $c_1(v_2) = c_1(v_1v_4) = r + 3$; the remaining vertices are colored with color 1.

Now we will verify that $W_4^{r,1}$ is total rainbow connected under this coloring. Let u and w be any two vertices of $W_4^{r,1}$. Clearly, there exists a total rainbow uw-path when u and w are adjacent. Thus, we assume that u and w are not adjacent in the sequel. We divide the remaining verification into the following cases according to the locations of u and w.

Case 1. $u, w \in \{v_1, v_2, v_3, v_4\}$. Then, uvw is a total rainbow uw-path.
Case 2. $u, w \in \{u_1, u_2, \ldots, u_r\}$. Then, uv_1w is a total rainbow uw-path.
Case 3. $u = w_1$.
Subcase 3.1 $w \in \{v, v_1, v_3\}$. Since $r \geq 4$, then uv_2w is a total rainbow uw-path.
Subcase 3.2 $w = v_4$. Then, uv_2vw is a total rainbow uw-path.
Subcase 3.3 $w = u_i$, where $i \in \{1, 2, \ldots, r\}$. Then, uv_2v_1w is a total rainbow uw-path if $i = 1$; uv_2v_1w is a total rainbow uw-path if $i = 2, 3, \ldots, r$.
Case 4 $u \in \{v_2, v_3, v_4, v\}$ and $w = u_i$, where $i \in \{1, 2, \ldots, r\}$.
Subcase 4.1 $u = v_2$. Then, uvv_1w is a total rainbow uw-path if $i = 1$, and uv_1w is a total rainbow uw-path if $i = 2, 3, \ldots, r$.
Subcase 4.2 $u = v_3$. Then, uvv_1w is a total rainbow uw-path if $i = 1, 4, 5, \ldots, r$, and uvv_2v_1w is a total rainbow uw-path if $i = 2$, and uv_2v_1w is a total rainbow uw-path if $i = 3$.
Subcase 4.3 $u = v_4$. Then, uv_1w is a total rainbow uw-path if $i = 1, 2, \ldots, r$.
Subcase 4.4 $u = v$. Then, uv_1w is a total rainbow uw-path if $i = 1, 3, 4, \ldots, r$, and uv_2v_1w is a total rainbow uw-path if $i = 2$.

By the above analysis, we get that $W_4^{r,1}$ is total rainbow connected under the coloring c_1. Hence, $trc(W_4^{r,1}) \leq r + 3$. Therefore, $trc(W_4^{r,1}) = r + 3$ for $r \geq 4$.

If $t = 7$ and $r \geq 5$, we give the graph $W_7^{r,1}$ a total coloring c_2 using $r + 3$ colors as follows:
$c_2(v_1u_i) = i$ for $1 \leq i \leq r$; $c_2(vv_4) = c_2(v_1v_2) = 1$; $c_2(vv_1) = c_2(v_5v_6) = c_2(v_3v_4) = 2$; $c_2(v) = 3$; $c_2(vv_2) = c_2(vv_7) = c_2(v_3) = c_2(v_6) = 4$; $c_2(vv_3) = c_2(v_1v_7) = c_2(v_4v_5) = 5$; $c_2(v_2w_1) = c_2(v_2v_3) = c_2(v_5) = c_2(v_7) = r + 1$; $c_2(v_1) = c_2(v_4) = c_2(vv_6) = r + 2$; $c_2(v_2) = c_2(vv_5) = c_2(v_6v_7) = r + 3$; the remaining vertices can be colored by arbitrary colors in $1, 2, \ldots, r + 3$.

Similar to the case $t = 4$, one can check that $W_7^{r,1}$ is total rainbow connected under the coloring c_2 when $r \geq 5$, which deduces that $trc(W_7^{r,1}) = r + 3$.

If $t = 10$ and $r \geq 8$, we define a total coloring c_3 for $W_{10}^{r,1}$ with $r+3$ colors as follows. $c_3(v_1u_i) = i$ for $1 \leq i \leq r$; $c_3(v_1v_{10}) = c_3(v_2v_3) = c_3(v_6v_7) = 1$; $c_3(vv_{10}) = c_3(v_3v_4) = c_3(v_9) = 2$; $c_3(v_4) = c_3(v_8) = 3$; $c_3(vv_6) = c_3(v_1v_2) = 4$; $c_3(vv_1) = c_3(v_5v_6) = c_3(v_8v_9) = 5$; $c_3(v) = 6$; $c_3(vv_2) = c_3(vv_8) = c_3(v_4v_5) = 7$; $c_3(v_3) = c_3(v_7v_8) = 8$; $c_3(v_2w_1) = c_3(vv_3) = c_3(v_5) = c_3(v_7) = c_3(v_{10}) = r + 1$; $c_3(v_1) = c_3(v_6) = c_3(vv_9) = r + 2$; $c_3(v_2) = c_3(vv_4) = c_3(vv_5) = c_3(vv_7) = c_3(v_9v_{10}) = r + 3$; the remaining vertices are colored by color 1.

We can easily verify that $W_{10}^{r,1}$ is total rainbow connected under the coloring c_3. Thus, $trc(W_{10}^{r,1}) \leq r + 3$, which implies $trc(W_{10}^{r,1}) = r + 3$ for $r \geq 8$.

The proof of the result is thus complete. ∎

Lemma 9. *Let r be an integer. Then, $strc(W_4^{r,1}) = r+4$ for $r \geq 4$; $strc(W_7^{r,1}) = r+5$ for $r \geq 5$; $strc(W_{10}^{r,1}) = r + 6$ for $r \geq 8$.*

Proof. Let c be a strong total rainbow coloring of $W_t^{r,1}$. Set $X = c(v_1) \cup c(v_2) \cup c(v_2w_1) \cup (\bigcup_{i=1}^{r} c(v_1u_i))$. We first present a claim here.

Claim 2. The colors $c(v_1), c(v_2), c(v_2w_1), c(v_1u_1), c(v_1u_2), \ldots, c(v_1u_r)$ are pairwise distinct, namely, $|X| = r + 3$.

Proof of Claim 2. For $1 \leq i \leq r$, the unique total rainbow w_1u_i-geodesic is $w_1v_2v_1u_i$, and thus, $c(v_1)$, $c(v_2)$, $c(v_2w_1)$ and $c(v_1u_i)$ are different. Additionally, for two vertices u_i and u_j, where $i \neq j$ and $i, j \in \{1, 2, \ldots, r\}$, the unique total rainbow u_iu_j-geodesic is $u_iv_1u_j$, and hence, $c(v_1u_i)$ and $c(v_1u_j)$ are distinct. To sum up, the claim holds.

If $t = 4$ and $r \geq 4$, for $1 \leq i \leq r$, the unique total rainbow w_1u_i-geodesic is $w_1v_2v_1u_i$, and thus, $c(v_1v_2)$ is different from colors $c(v_1), c(v_2), c(v_2w_1), c(v_1u_i)$, and hence, $c(v_1v_2) \notin X$, which implies $strc(W_4^{r,1}) \geq |X| + 1 = r + 4$. Now, we give the upper bound for $strc(W_4^{r,1})$ by defining a total coloring f_1 of $W_4^{r,1}$ as follows:
$f_1(v_1u_i) = i$ for $1 \leq i \leq r$; $f_1(v_3) = f_1(v_4) = f_1(v_2v_3) = f_1(v_3v_4) = 1$; $f_1(vv_2) = 2$; $f_1(vv_4) = 3$; $f_1(v_2w_1) = f_1(vv_3) = r + 1$; $f_1(v_1) = r + 2$; $f_1(v_1v_2) = f_1(v) = r + 3$; $f_1(vv_1) = f_1(v_1v_4) = f_1(v_2) = r + 4$; the remaining vertices are colored by color 1.

By simple and tedious checking, we can obtain that f_1 is a strong total rainbow coloring for $W_4^{r,1}$ with $r + 4$ colors, which deduces that $strc(W_4^{r,1}) \leq r + 4$. Therefore, $strc(W_4^{r,1}) = r + 4$ for $r \geq 4$.

If $t = 7$ and $r \geq 5$. First, we prove the lower bound for $strc(W_7^{r,1})$. In graph $W_7^{r,1}$, the unique total rainbow v_5w_1-geodesic is $v_5vv_2w_1$, and thus, $c(v)$ and $c(vv_5)$ are different from $c(v_2)$ and $c(v_2w_1)$. For $1 \leq i \leq r$, the unique total rainbow v_5u_i-geodesic is $v_5vv_1u_i$, and so, $c(v)$ and $c(vv_5)$ are also different from $c(v_1)$ and $c(v_1u_i)$. Therefore, $c(vv_5) \neq c(v)$ and $c(vv_5), c(v) \notin X$. Hence, $strc(W_7^{r,1}) \geq |X| + 2 = r + 5$.
Now, we prove the upper bound for $strc(W_7^{r,1})$ by defining a total coloring f_2 as follows:
$f_2(v_1u_i) = i$ for $1 \leq i \leq r$; $f_2(vv_6) = f_2(v_3v_4) = 1$; $f_2(vv_3) = f_2(v_3) = 2$; $f_2(vv_7) = f_2(v_2v_3) = 3$; $f_2(v_2w_1) = f_2(vv_1) = f_2(v_7) = r + 1$; $f_2(v_1) = f_2(vv_2) = r + 2$; $f_2(v_2) = f_2(vv_4) = f_2(v_1v_7) = r + 3$; $f_2(v_1v_2) = f_2(v_6v_7) = f_2(v) = r + 4$; $f_2(vv_5) = r + 5$; the remaining vertices and edges are colored by color 1.

It is easy to show that f_2 is a strong total rainbow coloring for $W_7^{r,1}$ with $r + 5$ colors, which implies $strc(W_7^{r,1}) \leq r + 5$. Combining the lower and upper bounds, we can arrive at $strc(W_7^{r,1}) = r + 5$ for $r \geq 5$.

If $t = 10$ and $r \geq 8$. Similar to the argument in the case that $t = 7$ and $r \geq 5$, the colors $c(v)$, $c(vv_5)$, $c(vv_6)$, $c(vv_7)$, $c(vv_8)$ are different from the colors in X. Since the unique total rainbow v_5v_8-geodesic is v_5vv_8, then $c(v)$, $c(vv_5)$ and $c(vv_8)$ are also distinct. Therefore, we can obtain $strc(W_{10}^{r,1}) \geq |X| + 3 = r + 6$. This is a lower bound for $strc(W_{10}^{r,1})$. In the following we will give an upper bound for $strc(W_{10}^{r,1})$ by defining a coloring f_3 as follows:
$f_3(v_1u_i) = i$ for $1 \leq i \leq r$; $f_3(vv_9) = f_3(vv_{10}) = f_3(v_2v_3) = 1$; $f_3(v_3) = 2$; $f_3(v_3v_4) = 3$; $f_3(v_2w_1) = f_3(vv_3) = f_3(vv_4) = f_3(v_9v_{10}) = r + 1$; $f_3(v_1) = f_3(vv_2) = r + 2$; $f_3(v_2) =$

$f_3(vv_1) = f_3(v_1v_{10}) = r+3$; $f_3(v) = f_3(v_{10}) = f_3(v_1v_2) = r+4$; $f_3(vv_7) = f_3(vv_8) = r+5$; $f_3(vv_5) = f_3(vv_6) = r+6$; the remaining vertices and edges are colored by color 1.

One can check that f_3 is a strong total rainbow coloring for $W_{10}^{r,1}$ using $r+6$ colors, and thus, $strc(W_{10}^{r,1}) \leq r+6$, which deduces $strc(W_{10}^{r,1}) = r+6$ for $r \geq 8$.

The proof is thus complete. ∎

Applying Lemmas 8 and 9, we can arrive at the following result directly.

Theorem 7. *Let a be an integer. Then, we have*
(i) If $a \geq 7$, there exists a connected graph G satisfying $trc(G) = a$ and $strc(G) = a+1$;
(ii) If $a \geq 8$, there exists a connected graph G satisfying $trc(G) = a$ and $strc(G) = a+2$;
(iii) If $a \geq 11$, there exists a connected graph G satisfying $trc(G) = a$ and $strc(G) = a+3$.

Proof. (i) Let $G = W_4^{a-3,1}$. Since $a \geq 7$, then $a-3 \geq 4$. Thus, by Lemmas 8 and 9, we get $trc(G) = trc(W_4^{a-3,1}) = a$ and $strc(G) = strc(W_4^{a-3,1}) = a+1$.

(ii) Let $G = W_7^{a-3,1}$. Since $a \geq 8$, then $a-3 \geq 5$. By Lemmas 8 and 9, one can obtain that $trc(G) = trc(W_7^{a-3,1}) = a$ and $strc(G) = strc(W_7^{a-3,1}) = a+2$.

(iii) Let $G = W_{10}^{a-3,1}$. Since $a \geq 11$, then $a-3 \geq 8$. Hence, by Lemmas 8 and 9, we have that $trc(G) = trc(W_{10}^{a-3,1}) = a$ and $strc(G) = strc(W_{10}^{a-3,1}) = a+3$.

The proof of this result is thus complete. ∎

Combing the Remark 1, Theorem 6 and Theorem 7, we can get the main result of this section as follows.

Theorem 8. *Let a and b be any two integers.*
(i) If $a = 5$ or $a \geq 11$, and $b \geq a$, then there exists a connected graph G such that $trc(G) = a$ and $strc(G) = b$;
(ii) If $a = 6$, and $b = 6$ or $b \geq 10$, then there exists a connected graph G such that $trc(G) = a$ and $strc(G) = b$;
(iii) If $a = 7$, and $b = 7, 8$ or $b \geq 11$, then there exists a connected graph G such that $trc(G) = a$ and $strc(G) = b$;
(iv) If $a = 8$, 9 or 10, and $a \leq b \leq a+2$ or $b \geq a+4$, then there exists a connected graph G such that $trc(G) = a$ and $strc(G) = b$.

Remark 2: For any pair of positive integers a and b, we can almost always construct a connected graph G such that $trc(G) = a$ meanwhile $strc(G) = b$. Except for four cases: (1) $a \leq 4$ and $b \geq a$; (2) $a = 6$ and $b = 7, 8, 9$; (3) $a = 7$ and $b = 9, 10$; (4) $a = 8, 9, 10$, and respectively, $b = 11, 12, 13$. These remaining cases will be studied in our future work.

5 Concluding remark

As mentioned earlier, $strc(G) \geq \max\{srvc(G), src(G)\}$ holds trivially for a nontrivial connected graph G. Moreover, if G is complete, then $strc(G) = 1 = src(G) = \max\{srvc(G), src(G)\}$, that is, the bound is sharp. On the other hand, since $strc(W_n) = \lceil \frac{n}{3} \rceil + 1$ for $n \geq 4$, while $srvc(W_n) = 1$ (see [13]), then $src(W_n) = \lceil \frac{n}{3} \rceil$ (see [4]). Then, there exists a graph G of order n such that $strc(G) > srvc(G)$ and $strc(G) > src(G)$ for any $n \geq 4$. In conclusion, for $srvc(G)$, we conjecture that $strc(G) > srvc(G)$. For $src(G)$, we at present cannot be sure that how tight is the inequality $strc(G) \geq src(G)$. Does there exist a graph G such that $strc(G) = src(G) = s$ for sufficiently large integer s ? or, can we possibly expect that $strc(G) > src(G)$ for any non-complete graph G ? These problems will be the objects of our future study.

References

[1] J.A. Bondy, U.S.R. Murty, *Graph Theory*, Springer, Berlin, 2008.

[2] Y. Caro, A. Lev, Y. Roditty, Zs. Tuza, R. Yuster, On rainbow connection, *Electron. J. Combin.* **15**(2008), R57.

[3] S. Chakraborty, E. Fischer, A. Matsliah, R. Yuster, Hardness and algorithms for rainbow connectivity, 26th International Symposium on Theoretical Aspects of Computer Science STACS 2009 (2009), 243–254. Also, see *J. Combin. Optim.* **21**(2011) 330–347.

[4] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang, Rainbow connection in graphs, *Math. Bohem.* **133**(2008) 85–98.

[5] L. Chen, X. Li, H. Lian, Further hardness results on the rainbow vertex connection number of graphs, *Theoret. Comput. Sci.* **481**(2013) 18–23.

[6] L. Chen, X. Li, Y. Shi, The complexity of determining the rainbow vertex-connection of a graph, *Theoret. Comput. Sci.* **412**(2011) 4531–4535.

[7] X. Chen, X. Li, A solution to a conjecture on the rainbow connection number, *Ars Combin.* **104**(2012) 193–196.

[8] X. Huang, X. Li, Y. Shi, Note on the hardness of rainbow connections for planar and line graphs, *Bull. Malays. Math. Sci. Soc.* **38(3)**(2015) 1235–1241.

[9] H. Jiang, X. Li, Y. Zhang, Upper bounds for the total rainbow connection of graphs, *J. Combin. Optim.*, DOI 10.1007/s10878-015-9874-x, in press.

[10] M. Krivelevich, R. Yuster, The rainbow connection of a graph is (at most) reciprocal to its minimum degree, *J. Graph Theory* **63**(2010) 185–191.

[11] H. Li, X. Li, S. Liu, The (strong) rainbow connection numbers of Cayley graphs on Abelian groups, *Comput. Math. Appl.* **62(11)**(2011) 4082–4088.

[12] S. Li, X. Li, Y. Shi, Note on the complexity of deciding the rainbow (vertex-) connectedness for bipartite graphs, *Appl. Math. Comput.* **258**(2015) 155–161.

[13] X. Li, Y. Mao, Y. Shi, The strong rainbow vertex-connection of graphs, *Util. Math.* **93**(2014) 213–223.

[14] X. Li, Y. Shi, On the rainbow vertex-connection, *Discuss. Math. Graph Theory* **33**(2013) 307–313.

[15] X. Li, Y. Shi, Y. Sun, Rainbow connections of graphs: A survey, *Graphs Combin.* **29**(2013) 1–38.

[16] X. Li, Y. Sun, *Rainbow Connections of Graphs*, Springer Briefs in Math., Springer, New York, 2012.

[17] X. Li, Y. Sun, On the strong rainbow connection of a graph, *Bull. Malays. Math. Sci. Soc.* **36(2)**(2013) 299–311.

[18] H. Liu, Â. Mestre, T. Sousa, Total rainbow k-connection in graphs, *Discrete Appl. Math.* **174**(2014) 92–101.

[19] Y. Ma, Total rainbow connection number and complementary graph, *Results Math.*, in press.